

PRODUCT INFORMATION

PANADOL OSTEO


1. NAME OF THE MEDICINE

Paracetamol

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Active ingredient: Paracetamol 665 mg

Chemical structure:

Active ingredients	Chemical structure	CAS Registry Number
Paracetamol	 <chem>CC(=O)Nc1ccc(O)cc1</chem>	103-90-2

Excipients:

For the full list of excipients, see section 6.1 List of Excipients.

3. PHARMACEUTICAL FORM

White to off-white film coated capsule shaped tablets with flat edges. Embossed with the logo "8" on the front face and plain on the back face.

4. CLINICAL PARTICULARS

4.1 Therapeutic Indications

PANADOL OSTEO provides effective relief from persistent pain for up to 8 hours. Effective for the relief of persistent pain associated with osteoarthritis and muscular aches and pains such as backache. Provides effective temporary relief of pain and discomfort associated with: headache, tension headache, cold and flu, period pain, toothache and pain after dental procedures. Reduces fever.

4.2 Dose and Method of Administration

Adults and children aged 12 years and over: 2 caplets swallowed whole three times a day every 6 to 8 hours. Do not chew or suck, as it impairs the sustained release properties. Maximum of 6 caplets in 24 hours.

Do not use for more than a few days at a time in adults except on medical advice.

Children under 12 years: Not recommended for children under the age of 12 years.

Should not be used for more than 48 hours for children aged 12 – 17 years except on medical advice.

Take with water or other fluid.

Can be taken with or without food.

Doses should be equally spaced throughout the day.

The caplets must not be crushed.

Do not exceed the stated dose.

The lowest dose necessary to achieve efficacy should be used for the shortest duration of treatment.

Should not be used with other paracetamol-containing products.

Minimum dosing interval: 6 hours.

Maximum daily dose: 4000 mg

4.3 Contraindications

Contraindicated in patients with a previous history of hypersensitivity to paracetamol or to any of the excipients.

4.4 Special warnings and precautions for use

Contains paracetamol. Do not use with any other paracetamol- containing products. The concomitant use with other products containing paracetamol may lead to an overdose.

Paracetamol overdose may cause liver failure which may require liver transplant or lead to death.

Underlying liver disease increases the risk of paracetamol-related liver damage. Patients who have been diagnosed with liver or kidney impairment must seek medical advice before taking this medication. The restrictions related to the use of paracetamol products in patients with hepatic impairment are primarily a consequence of the paracetamol content of the drug.

Cases of hepatic dysfunction/failure have been reported in patients with depleted glutathione levels, such as those who are severely malnourished, anorexic, have a low body mass index, are chronic heavy users of alcohol or have sepsis.

In patients with glutathione depleted states the use of paracetamol may increase the risk of metabolic acidosis.

If symptoms persist, medical advice must be sought.

Keep out of sight and reach of children.

Use in children

Not recommended for children under 12 years of age.

4.5 Interaction with other medicines and other forms of interaction

The following interactions with paracetamol have been noted:

The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol with increased risk of bleeding; occasional doses have no significant effect. Anticoagulant dosage may require reduction if paracetamol and anticoagulants are taken for a prolonged period of time.

Paracetamol absorption is increased by substances that increase gastric emptying, e.g. metoclopramide.

Paracetamol absorption is decreased by substances that decrease gastric emptying, e.g. propantheline, antidepressants with anticholinergic properties, and narcotic analgesics.

Paracetamol may increase chloramphenicol concentrations.

The risk of paracetamol toxicity may be increased in patients receiving other potentially hepatotoxic drugs or drugs that induce liver microsomal enzymes such as alcohol and anticonvulsant agents.

Paracetamol excretion may be affected and plasma concentrations altered when given with probenecid.

Colestyramine reduces the absorption of paracetamol if given within 1 hour of paracetamol.

4.6 Pregnancy and Lactation

Use in pregnancy

Category A

Paracetamol has been taken by a large number of pregnant women and women of childbearing age without any proven increase in the frequency of malformations or other direct or indirect harmful effects on the foetus having been observed.

As with the use of any medicine during pregnancy, pregnant women should seek medical advice before taking paracetamol. The lowest effective dose and shortest duration of treatment should be considered.

Use in Lactation

Paracetamol is excreted in breast milk. Human studies with paracetamol have not identified any risk to lactation or the breast-fed offspring. These results are based on immediate release preparations of paracetamol. There is no data available on the excretion of sustained-release paracetamol preparations in breast milk. However, it is not expected that PANADOL OSTEO would provide any increase in the excretion of paracetamol in breast milk as this product is designed to maintain rather than increase plasma paracetamol concentrations compared to immediate release preparations. Maternal ingestion of paracetamol in usual analgesic doses does not appear to present a risk to the breastfed infant.

4.7 Effects on ability to drive and use machines

The effects of this medicine on a person's ability to drive and use machines were not assessed as part of its registration.

4.8 Adverse effects

Adverse events from historical clinical trial data are both infrequent and from small patient exposure. Accordingly, events reported from extensive post-marketing experience at therapeutic/labelled doses and considered attributable are tabulated below by System Organ Class and frequency.

The following convention has been utilised for the classification of undesirable effects: very common ($\geq 1/10$), common ($\geq 1/100, < 1/10$), uncommon ($\geq 1/1,000, < 1/100$), rare ($\geq 1/10,000, < 1/1,000$), very rare ($< 1/10,000$), not known (cannot be estimated from available data).

Adverse event frequencies have been estimated from spontaneous reports received through post-marketing data.

Body System	Undesirable Effect	Frequency
Blood and lymphatic system disorders	Thrombocytopenia	Very rare
Immune system disorders	Anaphylaxis Cutaneous hypersensitivity reactions including, among others, skin rashes, angioedema, Stevens Johnson syndrome and Toxic Epidermal Necrolysis.	Very rare
Respiratory, thoracic and mediastinal disorders	Bronchospasm, especially in patients sensitive to aspirin and other NSAIDS	Very rare
Hepatobiliary disorders	Hepatic dysfunction	Very rare

4.9 Overdose

Poisons Information Centre

If an overdose is taken or suspected, contact the Poisons Information Centre immediately for advice (131 126), or the patient should go to the nearest hospital straight away. This should be done even if they feel well because of the risk of delayed, serious liver damage (see ADVERSE EFFECTS).

Because PANADOL OSTEO is a sustained-release formulation of paracetamol, absorption will be prolonged in overdose. It is recommended that for the management of overdose, where PANADOL OSTEO is suspected, that an additional plasma paracetamol level be obtained 4-6 hours after the initial measurement. If either level is above or close to the treatment line on the paracetamol overdose nomogram, administration of antidote would be indicated.

Treatment

Paracetamol overdose may cause liver failure which may require liver transplant or lead to death. Acute pancreatitis has been observed, usually with hepatic dysfunction and liver toxicity.

Immediate medical management is required in the event of an overdose, even if the symptoms of overdose are not present.

Administration of N-acetylcysteine may be required.

In cases of overdosage, methods of reducing absorption of ingested drug are important. Activated charcoal may reduce absorption of the medicine if given within one hour after oral ingestion. In patients who are not fully conscious or have impaired gag reflex, consideration should be given to administering activated charcoal via a nasogastric tube, once the airway is protected.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Paracetamol is a para-aminophenol derivative that exhibits analgesic and anti-pyretic activity. Its mechanism of action is believed to include inhibition of prostaglandin synthesis, primarily within the central nervous system. It does not possess anti-inflammatory activity. It provides relief from mild to moderate pain and fever.

The combination of immediate release and sustained release paracetamol provides pain relief, which may last up to 8 hours.

CLINICAL TRIALS

Chronic Pain

In patients with pain associated with osteoarthritis of the knee, PANADOL OSTEO (2 tablets taken three times daily) and standard immediate release paracetamol (2 tablets taken 4 times daily) were clinically equivalent at a total daily dose of 4 g based on patient global assessment after treatment for 7 days.

PANADOL OSTEО and standard immediate release paracetamol were not significantly different for a range of secondary efficacy parameters including pain during the day, pain on walking, pain relief, number of times woken during the night due to pain and duration of morning stiffness.

Since PANADOL OSTEО (three times daily) was clinically equivalent to standard immediate release paracetamol (four times daily), it was concluded that PANADOL OSTEО provides pain relief for up to 8 hours after dosing.

Acute Pain

In patients with post-surgical dental pain, a single dose of PANADOL OSTEО (2 tablets) was therapeutically equivalent to standard immediate release paracetamol (2 tablets) based on patient global assessment 4 hours after treatment. Onset of action was apparent 30 minutes after administration.

There was no significant difference between PANADOL OSTEО and standard immediate release paracetamol in either development of analgesia or peak analgesic effect. Trends in favour of PANADOL OSTEО were observed at the later time points. Furthermore, PANADOL OSTEО was significantly more effective than standard immediate release paracetamol for the summed pain intensity difference at 6 hours ($p = 0.0344$) and 8 hours ($p = 0.0500$), as measured on a visual analogue scale.

Summary

From these results, it was concluded that PANADOL OSTEО has a similar time to onset of action compared to standard immediate release paracetamol and provides more prolonged analgesia than standard immediate release paracetamol. For the patient, this translates to longer lasting pain relief and the improved convenience of fewer doses. This is as expected for a formulation containing sustained release paracetamol and consistent with results from the pharmacokinetic studies.

5.2 Pharmacokinetic properties

Absorption

Paracetamol is rapidly and almost completely absorbed from the gastrointestinal tract. Food intake delays paracetamol absorption.

PANADOL OSTEО is a unique bi-layer tablet incorporating an immediate release and a sustained release dose of paracetamol.

The sustained release layer contains HPMC polymer, which rapidly hydrates to form a gel layer at the matrix periphery. The drug is then released from the matrix by a combination of diffusion and erosion of the gel layer.

Bioequivalence

PANADOL OSTEО releases drug at a rate which ensures that therapeutically active plasma paracetamol concentrations are rapidly attained and maintained until up to 8 hours after administration.

PANADOL OSTEO PANADOL OSTEO and standard immediate release paracetamol were bioequivalent in volunteers with respect to dose-corrected $AUC_{(0-t)}$ and $AUC_{(0-inf)}$ in both the fed and fasted states following administration of a single dose. This indicates that the extent of paracetamol absorption from PANADOL OSTEO was equivalent to that of standard immediate release paracetamol. Food had little effect on the extent of paracetamol absorption from PANADOL OSTEO demonstrating that PANADOL OSTEO is suitable to be taken with or without meals. Paracetamol was rapidly absorbed after administration of PANADOL OSTEO and was generally measurable in plasma within 15 minutes in fasted subjects. Mean plasma paracetamol concentrations above the minimum level required for analgesia ($>4\text{mcg/mL}$) were maintained until up to 6 to 7 hours after administration in fasted subjects and 7 to 8 hours in fed subjects.

At steady state, PANADOL OSTEO was bioequivalent with standard immediate release paracetamol based on the comparison of AUCs during the final 24 hour dosing period of the study. Furthermore, comparison of the pharmacokinetic parameters indicated that PANADOL OSTEO has the characteristics of a formulation containing sustained release paracetamol.

Fluctuations in the peak and trough values for plasma paracetamol concentrations were significantly smaller for PANADOL OSTEO than for standard immediate release paracetamol (mean fluctuation index = 0.957 and 1.388, respectively, $p < 0.001$). Consequently, PANADOL OSTEO provided more consistent levels of paracetamol. Furthermore, the AUCs at steady state were equivalent indicating that there was no additional accumulation of paracetamol from PANADOL OSTEO compared to standard immediate release paracetamol.

Distribution

Paracetamol is distributed into most body tissues. Binding to the plasma proteins is minimal at therapeutic concentrations but increases with increasing doses.

Metabolism

Paracetamol is metabolised extensively in the liver and excreted in the urine mainly as inactive glucuronide and sulphate conjugates.

The metabolites of paracetamol include a minor hydroxylated intermediate which has hepatotoxic activity. This intermediate metabolite is detoxified by conjugation with glutathione. However, it can accumulate following paracetamol overdose (more than 150 mg/kg or 10 g total paracetamol ingested) and, if left untreated, can cause irreversible liver damage.

Paracetamol is metabolised differently by premature infants, newborns, infants and young children compared to adults, the sulfate conjugate being predominant.¹

¹ Core Paracetamol PI

Excretion

Paracetamol is excreted in the urine mainly as the glucuronide and sulfate conjugates. Less than 5% is excreted unchanged. Approximately 85% of a dose of paracetamol is excreted in urine as free and conjugated paracetamol within 24 hours of ingestion. Administration of paracetamol to patients with moderate to severe renal impairment may result in accumulation of paracetamol conjugates.² The elimination half-life varies from one to three hours.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Hypromellose, pregelatinised maize starch, Povidone, Croscarmellose sodium, Magnesium stearate, Stearic acid, Glycerol Triacetate, Carnauba wax.

Contains no sugar, lactose or gluten.

6.2 Incompatibilities

Incompatibilities were either not assessed or not identified as part of the registration of this medicine.

6.3 Shelf life

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

6.4 Special precautions for storage

Store below 30°C.

6.5 Nature and contents of container

Packs of 6, 12, 18, 24, 36, 48 and 96 caplets.

Bottles 96, 400 and 1000 caplets.

Not all pack/ bottle sizes may be marketed.

7. MEDICINE SCHEDULE

S3 - Pharmacist Only Medicine

8. SPONSOR

Haleon Australia Pty Ltd
Level 48, 8 Parramatta Square
10 Darcy Street, Parramatta NSW 2150

FREECALL Australia: 1800 028 533

Website: www.haleon.com

² American Hospital Formulary Service Drug Information 2012

9. DATE OF FIRST APPROVAL

Blister pack: 24/09/2015

10. DATE OF REVISION OF TEXT

31 May 2024

Summary table of changes

<u>Section changes</u>	<u>Summary of new changes</u>
8	Update sponsor details from GSK to Haleon

Trademarks are owned by or licensed to the Haleon group of companies.