PENTASA® suppositories

Consumer Medicine Information (CMI) summary

The <u>full CMI</u> on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using PENTASA?

PENTASA contains the active ingredient mesalazine. PENTASA is used to treat proctitis: a disease with inflammation, ulcers and sores in the rectum (back passage) causing bleeding, stomach pain, and diarrhoea.

For more information, see Section 1. Why am I using PENTASA? in the full CMI, starting from page 2.

2. What should I know before I use PENTASA?

Do not use PENTASA if you have ever had an allergic reaction to mesalazine or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section 2. What should I know before I use PENTASA? in the full CMI, starting from page 2..

3. What if I am taking other medicines?

Care may be needed when using PENTASA with some other medicines.

A list of these medicines is in Section 3. What if I am taking other medicines? in the full CMI, starting from page 2..

4. How do I use PENTASA?

- Use as directed by your doctor.
- Insert one suppository (1 g) into the rectum at bedtime.
- More instructions can be found in Section <u>4. How do I use PENTASA?</u> in the full CMI, starting from page 2. and in the detailed instructions in the pack.

5. What should I know while using PENTASA?

Things you should do	Remind any doctor, dentist or pharmasist
	dentist or pharmacist

	you visit that you are using PENTASA Have all blood tests recommended by your doctor If you are going to have surgery, tell the surgeon or anaesthetist that you are using this medicine If you become pregnant while using this medicine, tell your doctor immediately.
Things you should not do	 Do not stop using this medicine or lower the dosage suddenly without checking with your doctor Do not use PENTASA to treat any other complaints unless your doctor tells you to Do not give your medicine to anyone else, even if they have the same condition as you.
Driving or using machines	PENTASA is not expected to affect your

	ability to drive a car or operate machinery.
Looking after your medicine	 Keep your PENTASA in a cool dry place where the temperature stays below 25°C.

For more information, see Section <u>5. What should I know while using PENTASA?</u> in the full CMI, starting from page 2.

6. Are there any side effects?

All medicines can have side effects. Side effects reported from PENTASA can be minor and temporary. However, some side effects may need medical attention.

For more information, including what to do if you have any side effects, see Section <u>6. Are there any side</u> <u>effects?</u> in the full CMI, starting from page 2.

PENTASA® suppositories

Active ingredient(s): mesalazine

Consumer Medicine Information (CMI)

This leaflet provides important information about using PENTASA. You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using PENTASA.

Where to find information in this leaflet:

- 1. Why am I using PENTASA?
- 2. What should I know before I use PENTASA?
- 3. What if I am taking other medicines?
- 4. How do I use PENTASA?
- 5. What should I know while using PENTASA?
- 6. Are there any side effects?
- 7. Product details

1. Why am I using PENTASA?

PENTASA contains the active ingredient mesalazine.

PENTASA is an anti-inflammatory agent, similar to aspirin and is thought to reduce inflammation in the rectum.

PENTASA is used to treat proctitis: a disease with inflammation, ulcers and sores in the rectum (back passage) causing bleeding, stomach pain, and diarrhoea.

2. What should I know before I use PENTASA?

Warnings

Do not use PENTASA if:

- you are allergic to mesalazine, or any of the ingredients listed at the end of this leaflet always check the ingredients to make sure you can use this medicine
- have a severe kidney or liver problem.

Check with your doctor if you:

- have a known allergy to PENTASA, sulfasalazine or aspirin-like medicines, or any of the ingredients listed at the end of this leaflet
- take any medicines for any other condition
- have a kidney or liver problem
- have a bleeding disorder
- have a history of asthma.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section 6. Are there any side effects?

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

Your doctor can discuss with you the risks and benefits involved. PENTASA should be used with caution during pregnancy and breast-feeding and only if the potential benefits outweigh the possible risks in the opinion of the doctor. The underlying condition itself (inflammatory bowel disease) may increase risks for the pregnancy outcome.

Children

 Do not give PENTASA to a child 12 years old or under.

The safety and effectiveness of PENTASA in this age group have not been established.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription

from your pharmacy, supermarket or health food shop.

There is no information available on interactions between PENTASA and other medicines. However, mesalazine belongs to a group of medicines called salicylates that may interfere with some medicines.

Medicines that may have their effects increased if used with PENTASA include:

- anti-coagulants which are used to thin blood or stop blood clots (e.g. warfarin). When PENTASA is used with these medicines, it may increase the medicine's effects on thinning your blood.
- sulphonylureas which are used to lower blood sugar and treat diabetes (e.g. glibenclamide, glipizide).
 When PENTASA is used with these medicines, it may increase the medicine's effect on lowering your blood glucose.
- methotrexate which is used to treat some kinds of cancers, some skin conditions, and rheumatoid arthritis. When PENTASA is used with this medicine, it may increase the possibility of the medicine producing toxic effects on your body.
- glucocorticoids which are used to treat inflammation, swelling or allergies (e.g. prednisolone). When PENTASA is used with these medicines, it may increase the undesirable gastric effects caused by these medicines.

Medicines that may have their effects decreased if used with PENTASA include:

- probenicid or sulfinpyrazone which are used to treat gout and other conditions. When PENTASA is used with these medicines it may decrease the effectiveness of the medicine.
- spirinolactone or frusemide, which are used to increase the amount of urine produced and to lower blood pressure. When PENTASA is used with these medicines it may decrease the effectiveness of the medicine.
- rifampicin which is used to treat tuberculosis. When PENTASA is used with this medicine it may decrease the effectiveness of the medicine.

Medicines that may have a greater likelihood of reducing bone marrow activity if used with PENTASA include:

- azathioprine which is used to suppress the immune system
- mercaptopurine and thioguanine, used to treat leukaemia.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect PENTASA.

4. How do I use PENTASA?

This medicine helps to control your condition, but does not cure it. It is important to keep using your medicine even if you feel well.

How much to use

- The recommended dose is one suppository (1 g) into the rectum, once daily at bedtime
- Your doctor will prescribe a dosage regimen based on your individual needs, and this may differ from this recommendation
- Follow the instructions provided and use PENTASA until your doctor tells you to stop.

When to use PENTASA

PENTASA should be used at bedtime.

How to use PENTASA

- If possible, go to the toilet and empty your bowels before using a suppository. The suppository will work better if the bowel has been emptied
- Insert the contents of the suppository in the rectum (back passage)
- Detailed instructions for use are available in the pack
- Do not swallow the suppository.



Scan the QR code with your smartphone or tablet to access a video on administering PENTASA® Enema. Select the PENTASA® link on the website, enter the AUST R number from the PENTASA® box or Section 7 of this document, and scroll to resources. Alternatively, visit patients.ferring.com.au/gastroenterology/Access available only within Australia.

If you forget to use PENTASA

PENTASA should be used regularly at the same time each day.

If it is almost time for your next suppository, skip the dose you missed and use your next suppository when you are meant to.

Do not use a double dose to make up for the dose you missed.

If you use too much PENTASA or swallow the medicine:

If you think that you have used too much PENTASA, or if the medicine has been taken by mouth, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (by calling 13 11 26), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while using PENTASA?

Things you should do

Have all blood tests recommended by your doctor:

- PENTASA may cause kidney, liver or blood problems in a few people. You should have regular blood tests to check your kidney function
- Kidney stones may develop while using PENTASA.
 Symptoms may include pain in the sides of the abdomen and blood in the urine
- Take care to drink plenty of fluids while you are being treated with PENTASA.

If you are going to have surgery:

 Tell the surgeon or anaesthetist that you are using this medicine. It may affect other medicines used during surgery.

If you become pregnant while using this medicine, tell your doctor immediately.

Remind any doctor, dentist or pharmacist you visit that you are using PENTASA.

Things you should not do

- Do not stop using this medicine or lower the dosage suddenly without checking with your doctor. If you stop taking it suddenly, your condition may worsen
- Do not use PENTASA to treat any other complaints unless your doctor tells you to
- Do not give your medicine to anyone else, even if they have the same condition as you.

Driving or using machines

PENTASA is not expected to affect your ability to drive a car or operate machinery.

Looking after your medicine

Follow the instructions in the carton on how to take care of your medicine properly.

- Keep your PENTASA in the packaging until it is time to use your next dose.
- If you take PENTASA out of the packaging it may not keep well.
- Keep your PENTASA in a cool dry place (temperature stays below 25°C), away from moisture, heat or sunlight.
- Do not store it:
- in the bathroom or near a sink, or
- in the car or on window sills.

Keep it where children cannot reach it.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Side effects	What to do
Common side effects (affect more than 1 in 100 users):	These side effects are not usually serious but can become serious
 headache diarrhoea abdominal or stomach pain nausea vomiting 	Speak to your doctor if you have any of these side effects

Side effects	What to do
 flatulence (stomach discomfort or fullness, relieved by passing wind) 	
mild skin rash	
 discomfort, pain or itching in the anus or rectum, or feeling that you need to have a bowel movement, during or after inserting the medicine 	
Rare side effects (affect	
less than 1 in 1,000	
users):	
dizziness	
 photosensitivity (especially if you have eczema or dermatitis) 	
Very rare side effects (affect less than 1 in 10,000 users):	
 changes in blood counts e.g. a decrease in the number of red/white blood cells, and/or platelets 	

Side effects	What to do
 difficulty breathing, coughing (allergic and fibrotic lung reactions) changes in liver function e.g. raised liver enzymes hair loss (reversible) painful or swollen joints (arthralgia) muscle aches and pains (myalgia) disease affecting the skin, joints and kidneys with symptoms such as joint pain, fever and skin rash (lupus erythematosus-like reactions) low sperm count (reversible). drug fever e.g. fever-like symptoms such as chills 	
and aches Frequency unknown:	
• kidney stones	

Serious side effects

Serious side effects	What to do
Rare side effects (affect less than 1 in 1,000 users):	Call your doctor straight away, if you notice any of these serious side
 chest pain, irregular heartbeat, shortness of breath (myocarditis and pericarditis) 	effects.
 inflammation of the pancreas e.g. severe upper stomach pain, nausea and vomiting (increased amylase, acute pancreatitis) 	
Very rare side effects (affect less than 1 in 10,000 users):	
 bruising easily, unusual bleeding (e.g. nosebleeds), signs of infection such as fever, chills, sore throat and mouth ulcers (aplastic anemia or agranulocytosis) 	
 numbness, tingling or weakness of the arms 	

Serious side effects	What to do
and legs (peripheral neuropathy)	
 worsening of your condition e.g. severe diarrhea, bloody stools (pancolitis) 	
 liver disease e.g. yellowing of the skin or eyes, urine discolouration. 	
 kidney disease e.g. changes in urine output and/or discolouration, blood in urine and/or swelling 	
 red lesion or rash (erythema multiforme or dermatitis allergic) 	
Frequency unknown:	
 increased pressure in the brain, strong or frequent headaches, changes in vision, or ringing in the ears (idiopathic intracranial hypertension) 	

Serious side effects

What to do

Allergic reaction (very rare - affect less than 1 in 10,000 users):

Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects

 Sudden signs of allergic reactions such as rash, itching or hives, shortness of breath, wheezing, coughing, or swelling of limbs, face, lips, mouth, tongue or throat which may cause difficulty swallowing or breathing

Severe Cutaneous
Adverse Reactions
(SCAR) such as
Drug Reaction with
Eosinophilia and
Systemic Symptoms
(DRESS) (very rare affect less than 1 in
10,000 users); StevensJohnson Syndrome
(SJS) /Toxic epidermal
necrolysis (TEN)
(frequency unknown):

 Nausea, severe stomach cramps and/or pain, bloody diarrhoea,

Serious side effects	What to do
fever, facial swelling, severe headache and skin rash with severe blisters and bleeding of the eyes, mouth, lips, nose and genitals.	

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop using any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What PENTASA contains

Active ingredient (main ingredient)	mesalazine
Other ingredients (inactive ingredients)	povidone macrogol 6000 magnesium stearate
	talc
Potential allergens	This medicine does not contain sucrose, gluten, tartrazine or any other azo dyes.

Do not use this medicine if you are allergic to any of these ingredients.

What PENTASA looks like

PENTASA Suppositories are white to tan spotted, oblong shaped. Each suppository is protected in an aluminium foil blister. PENTASA Suppositories may be supplied in packs of 5, 10, 28 and 30. Not all pack sizes are distributed in Australia.

(AUST R 98034)

Who distributes PENTASA

PENTASA Suppositories are supplied in Australia by:

Ferring Pharmaceuticals Pty Ltd

Suite 2, Level 1, Building 1

20 Bridge Street

Pymble, NSW 2073, Australia.

PENTASA® is a registered trademark of Ferring B.V.

® = Registered trademark

This leaflet was prepared in October 2025.

AU-PA-2100040_v3.0