

# HUMULIN<sup>®</sup> R/HUMULIN<sup>®</sup> NPH/HUMULIN<sup>®</sup> 30/70 cartridges

## Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

### 1. Why am I using HUMULIN?

HUMULIN contains the active ingredient **biosynthetic human insulin (rbe)**. HUMULIN is used to **reduce high blood sugar (glucose) levels** in patients with diabetes. For more information, see Section [1. Why am I using HUMULIN?](#) in the full CMI.

### 2. What should I know before I use HUMULIN?

Do not use if you have ever had an allergic reaction to biosynthetic human insulin or any of the ingredients listed at the end of the CMI or when you have hypoglycaemia (low blood sugar).

**Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.**

For more information, see Section [2. What should I know before I use HUMULIN?](#) in the full CMI.

### 3. What if I am taking other medicines?

Some medicines may interfere with HUMULIN and affect how it works. A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

### 4. How do I use HUMULIN?

- Your doctor will tell you what dose of HUMULIN you need to use each day.
- HUMULIN cartridges are only for use with the HumaPen.

More instructions can be found in Section [4. How do I use HUMULIN?](#) in the full CMI.

### 5. What should I know while using HUMULIN?

<b>Things you should do</b>	<ul style="list-style-type: none"><li>• Remind any doctor, dentist, or pharmacist you visit that you are using HUMULIN.</li><li>• Monitor your blood sugar levels regularly.</li><li>• Tell your doctor if you become pregnant or start breastfeeding while using HUMULIN insulin.</li></ul>
<b>Things you should not do</b>	<ul style="list-style-type: none"><li>• Do not stop using HUMULIN without first checking with your doctor.</li><li>• Do not change your dose of HUMULIN without first checking with your doctor.</li><li>• Do not use HUMULIN when you have hypoglycaemia (low blood sugar) as it will make it worse.</li></ul>
<b>Driving or using machines</b>	<ul style="list-style-type: none"><li>• Be careful before you drive or use any machines or tools until you know how HUMULIN affects you.</li><li>• Hypoglycaemia (low blood sugar) may impair your ability to concentrate or react while driving. Do not drive if you feel like your blood sugar levels are low.</li></ul>
<b>Drinking alcohol</b>	<ul style="list-style-type: none"><li>• Tell your doctor if you drink alcohol. Alcohol may mask the symptoms of hypoglycaemia (low blood sugar).</li></ul>
<b>Looking after your medicine</b>	<ul style="list-style-type: none"><li>• Keep your cartridges in a fridge between 2°C - 8°C. Do not freeze.</li><li>• After first use you may keep the cartridges below 30°C for up to 21 days. Dispose after 21 days even if the cartridge contains insulin.</li></ul>

For more information, see Section [5. What should I know while using HUMULIN?](#) in the full CMI.

### 6. Are there any side effects?

The most common (and potentially the most severe) side effect of HUMULIN is hypoglycaemia (low blood sugar).

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

# HUMULIN<sup>®</sup> R/HUMULIN<sup>®</sup> NPH/HUMULIN<sup>®</sup> 30/70 cartridges

Active ingredient: biosynthetic human insulin (rbe)

## Consumer Medicine Information (CMI)

This leaflet provides important information about using HUMULIN. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using HUMULIN.**

Where to find information in this leaflet:

1. [Why am I using HUMULIN?](#)
2. [What should I know before I use HUMULIN?](#)
3. [What if I am taking other medicines?](#)
4. [How do I use HUMULIN?](#)
5. [What should I know while using HUMULIN?](#)
6. [Are there any side effects?](#)
7. [Product details](#)

## 1. Why am I using HUMULIN?

HUMULIN contains the active ingredient **biosynthetic human insulin (rbe)**. It is identical to the insulin produced by the pancreas.

There are 3 different types of HUMULIN cartridges:

- HUMULIN R
- HUMULIN NPH
- HUMULIN 30/70.

They all contain the same active ingredient but **HUMULIN R** is a solution, **HUMULIN NPH** is a suspension and **HUMULIN 30/70** is a pre-mixed suspension. Each will vary with how long they work in your body.

HUMULIN is used to **reduce high blood sugar (glucose) levels** in patients with diabetes.

Diabetes is a condition in which your pancreas does not produce enough insulin to control your blood sugar level. Extra insulin is therefore needed.

There are two types of diabetes:

Type 1 diabetes - patients with type 1 diabetes always need insulin to control their blood sugar levels.

Type 2 diabetes - some patients with type 2 diabetes may also need insulin after initial treatment with diet, exercise, and tablets.

## 2. What should I know before I use HUMULIN?

### Warnings

#### Do not use HUMULIN:

- if you are allergic to biosynthetic human insulin, or any of the ingredients listed at the end of this leaflet. Always check the ingredients to make sure you can use this medicine.

- when you have hypoglycaemia (low blood sugar).
- HUMULIN cartridges should not be used in the muscle (intramuscularly) or in the vein (intravenously).

#### Check with your doctor if you:

- have previously had an allergic reaction to any medicine used to treat your diabetes.
- are using, or have used, another type or brand of insulin. This may affect the dose of HUMULIN that you are prescribed.
- have or have had overactive thyroid glands (hyperthyroidism), overactive adrenal glands (Cushing's syndrome), enlargement of the face, hands, and feet (acromegaly), or a tumour of the adrenal gland (phaeochromocytoma) as these conditions may affect the dose of HUMULIN that you are prescribed.
- have or have had kidney or liver disease. This may affect the dose of HUMULIN that you are prescribed.
- have or have had heart disease and are also taking thiazolidinediones a type of medicine used to treat type 2 diabetes (such as pioglitazone, rosiglitazone). You may be more at risk to develop fluid retention (oedema) especially in the hands, ankles, or feet, or heart failure while using HUMULIN.
- are taking any other medications.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

### Pregnancy and breastfeeding

**Check with your doctor if you are pregnant or intend to become pregnant.** Pregnancy may make managing diabetes more difficult. Insulin needs usually decrease during the first three months of pregnancy and increase during the last six months.

**Talk to your doctor if you are breastfeeding or intend to breastfeed.** Your doctor may need to change the dose of HUMULIN while you are breastfeeding.

### Children (under 18 years)

The use of HUMULIN has not been studied in children under the age of 18.

## 3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins, or supplements that you buy without a prescription from your pharmacy, supermarket, or health food shop.

**You may need a lower dose of HUMULIN if you are also taking any of the following medicines:**

- anabolic steroids

- other medicines that treat diabetes
- aspirin or other salicylates
- a group of antibiotics, known as sulfonamides, used to treat infections
- certain medicines used to treat depression, such as monoamine oxidase inhibitors
- certain medicines used to treat high blood pressure or other heart conditions, such as guanethidine, captopril, enalapril, angiotensin II receptor blockers, and beta blockers
- certain medicines used to treat people with growth disorders and tumours, such as octreotide.

You may need a **higher dose of HUMULIN** if you are also taking any of the following medicines:

- medicines used for oral contraception ('the pill')
- medicines that affect your metabolism (corticosteroids such as prednisone and cortisone, and growth hormones)
- glucagon
- adrenaline
- medicines used in thyroid replacement therapy
- certain antibiotics used to treat tuberculosis, such as isoniazid
- certain medicines used to treat asthma, such as salbutamol or terbutaline
- certain medicines used to treat schizophrenia, bipolar disorder, or nausea and vomiting, such as phenothiazines
- certain medicines used to treat epilepsy, such as phenytoin.

You may need to **change (increase or decrease) the dose of HUMULIN** if you are also taking the following medicine:

- diuretics, a group of medicines that help remove excessive fluids in the body.

**Check with your doctor or pharmacist if you are not sure about what medicines, vitamins, or supplements you are taking and if these affect HUMULIN.**

## 4. How do I use HUMULIN?

### How much to use

Your doctor will tell you what dose of HUMULIN you need to use each day.

### When to use HUMULIN

Your doctor will tell you when to use HUMULIN.

### How to use HUMULIN in a re-useable pen

**HUMULIN cartridges are only for use with the HumaPen.**

The manufacturer's instructions for the HumaPen delivery systems should be followed for loading the cartridge, attaching the needle, and administering the insulin injection.

### Check your cartridge

1. Check the cartridge before using. Make sure you are using the correct cartridge. The solution of **HUMULIN**

**R** should be clear and colourless. The suspension in **HUMULIN NPH** and **HUMULIN 30/70** cartridges should be white and cloudy or milky.

### Prepare dose

1. Wash your hands.
2. **HUMULIN NPH** and **HUMULIN 30/70** must be rolled between the palms ten times and inverted ten times until the contents are uniformly mixed. **Do not** shake as this may cause frothing which may interfere with the correct measurement of the dose.
3. Inspect the cartridge for uniform mixing and repeat the above steps if necessary.
4. Put the cartridge into the pen.
5. Attach a needle. Remove the outer needle shield. Do not throw it away.
6. Pull off the inner needle shield and throw it away.

### Checking for insulin flow (Priming)

**Always prime your pen before injecting a dose.**

Priming may highlight a malfunction with the pen or needle. It also removes any air bubbles and ensures that you receive the correct dose.

1. Set the dose to **2 units**.
2. Hold the pen with the needle pointing up and tap the side of the pen so that any bubbles float to the top.
3. With the pen still pointing up, press the injection button. Priming is complete when a stream of insulin comes out of the needle. This is very important to ensure the correct dose is injected.
4. If a drop of insulin still does not appear, hold the pen with the needle facing upwards and away from your face. Attach a new needle. Repeat steps 1 - 3 again. If you are still unable to get insulin flowing out of the needle, do not use the pen. Return the pen to your doctor, diabetes educator or pharmacist to obtain a replacement pen.
5. Only after seeing a stream of insulin at the needle tip should you proceed to dial up your required dose. There may still be some small air bubbles left in the cartridge. These are harmless.
6. If the air bubble is too big, you should prime the pen again.

### Injecting dose

HUMULIN should be injected under the skin (subcutaneously). Your doctor or diabetes educator will show you how to properly inject HUMULIN.

**Under no circumstances should HUMULIN NPH or HUMULIN 30/70 be injected intravenously (directly into your veins).**

1. Dial your prescribed dose
2. Choose a site for injection (upper arms, thighs, buttocks, or abdomen). **Use of injection sites should be rotated** so that the same spot is not used more than once a month.
3. Stabilise the skin by spreading it or gently pinching up a large area. Insert the needle as instructed by your doctor.
4. Press the knob. Slowly count to 5 before pulling the needle out.

### **After injecting**

1. Release the skin and apply gentle pressure over the injection site for several seconds. Do not rub the area.
2. Using the outer needle cap, unscrew the needle and dispose of it safely into a sharps container. **Do not re-use the needle.**
3. Leave the cartridge in the pen until it needs to be replaced.

### **Further injections**

- Stop using the cartridge if the leading edge of the plunger has gone to, or beyond, the left side of the gauge on the side of the cartridge.
- You can see how much insulin is left by looking at the gauge on the side of the cartridge. The scale on the cartridge shows how many units you have left. If there is not enough for your dose, change the cartridge.
- **Do not mix any other insulin in your cartridge. Once the cartridge is empty, do not use it again.**

### **If you forget to use HUMULIN**

#### **Test your blood sugar level as soon as possible.**

If you use less HUMULIN insulin than you need, a high blood sugar level (hyperglycaemia) may occur.

Use it as soon as you remember, and then go back to using it as you would normally. **Do not use a double dose of HUMULIN.**

If you are unsure, talk to your doctor, diabetes educator, or pharmacist.

### **If you use too much HUMULIN**

Hypoglycaemia (low blood sugar) may occur if you have used more HUMULIN than you need. Check your blood sugar levels.

If your blood sugar is low (mild hypoglycaemia) eat glucose tablets, sugar or drink a sugary drink. Then eat fruit, biscuits, or a sandwich. This will often resolve mild hypoglycaemia or a minor insulin overdose.

If you get worse and your breathing is shallow and your skin gets pale, you may need urgent medical attention.

#### **You should immediately:**

- phone the Poisons Information Centre (**by calling Australia 13 11 26 / New Zealand: 0800 764 766**), or
- contact your doctor, or
- go to the Emergency department at your nearest hospital.

**You should do this even if there are no signs of discomfort or poisoning.**

## **5. What should I know while using HUMULIN?**

### **Things you should do**

- Remind any doctor, dentist, pharmacist, and diabetes educator you visit that you are using HUMULIN.
- Monitor your blood sugar levels regularly to ensure that your dose of HUMULIN is appropriate. More

frequent monitoring is recommended when changing insulin dose or type, or when you start taking a new medicine.

- Always carry a supply of HUMULIN cartridges and your pen.
- Always carry something to show you have diabetes.
- Always carry glucose or sugary foods in case you develop hypoglycaemia (low blood sugar).

### **Tell your doctor**

- if you become pregnant or start breastfeeding while using HUMULIN.
- if you notice changes to your eyesight. This may be a sign that your diabetes is not being controlled appropriately.
- Tell your doctor, pharmacist, or diabetes educator if you plan to travel. You may need to carry a letter explaining why you are taking injecting devices with you. Your doctor, pharmacist, or diabetes educator can provide you with some helpful information.

### **Things you should not do**

- Do not use HUMULIN when you have hypoglycaemia (low blood sugar) as it will make it worse.
- Do not stop using HUMULIN without first checking with your doctor.
- Do not change your insulin (dose or type) without first checking with your doctor.
- Do not give your HUMULIN to anyone else, even if they have the same condition as you. Your doctor has prescribed HUMULIN specifically for you.
- **Do not share needles, cartridges, or pens.**
- **Do not re-use needles or syringes.**
- Do not store your insulin pen with the needle attached. This will ensure that your insulin will not become contaminated, and prevent leakage, the re-entry of air, and potential needle clogs.
- Do not use HUMULIN if the packaging is torn or shows signs of tampering.
- Do not use **HUMULIN R** if the solution is cloudy, thickened, coloured, or has solid pieces in it, as it may no longer be good to use.
- Do not use **HUMULIN NPH** or **HUMULIN 30/70** if the suspension is no longer cloudy white after gentle shaking or if it has solid pieces in it, as it may no longer be good to use.
- Do not use HUMULIN if the cartridge appears frosted, as it may no longer be good to use.
- Do not use HUMULIN if you think it has been frozen or exposed to excessive heat. It will not work as well.

### **Hypoglycaemia (low blood sugar)**

Your blood sugar level may become too low if you:

- use too much HUMULIN
- exercise too much or excessively
- delay eating meals or snacks
- eat too little food
- are ill.

Symptoms of **mild to moderate** hypoglycaemia can come on suddenly and may include:

- weakness, trembling or shaking
- sweating
- light headedness/headache
- lack of concentration
- behaviour change
- dizziness
- tearfulness/crying
- irritability
- numbness around the lips/fingers
- hunger

**Tell your doctor if you have trouble recognising these symptoms, if you often have hypoglycaemia or if you have ever become unconscious after using insulin.** More frequent monitoring of your blood sugar level may be recommended, or your dose of HUMULIN may need to be changed.

Recognising these mild to moderate symptoms early may allow you to take the necessary steps to avoid more serious hypoglycaemia.

**Always carry some sugary food or drink with you.**

If you experience symptoms of mild to moderate hypoglycaemia, immediately eat some sugary food or drink, such as jelly beans, fruit juice or glucose tablets.

**If you do not feel better after eating/drinking some sugary food or drink, contact your doctor or go to the Emergency department at your nearest hospital.**

**Tell your relatives, friends, close workmates, or carers that you have diabetes.**

It is important that they also recognise the signs and symptoms of hypoglycaemia.

Make sure they know to give you some sugary food or drink for mild to moderate symptoms of hypoglycaemia.

**Severe** symptoms of hypoglycaemia may include:

- disorientation
- seizures, fits, or convulsions
- loss of consciousness.

**If you lose consciousness, make sure your relatives, friends, close workmates, or carers know:**

- to turn you on your side and **get medical help immediately.**
- not to give you anything to eat or drink to avoid choking.

Severe hypoglycaemia can lead to unconsciousness and in extreme cases death if not treated.

An injection of the hormone glucagon may speed up recovery from unconsciousness. This can be given by a relative, friend, workmate, or carer who knows how to give it.

If glucagon is used, have some sugary food or drink as soon as you are conscious again.

### **Hyperglycaemia (high blood sugar)**

Your blood sugar levels may become too high if you:

- miss doses of HUMULIN or use less than you need
- have uncontrolled diabetes
- exercise less than usual
- eat more carbohydrates than usual
- are ill or stressed.

**Contact your doctor if your blood sugar level is too high and you experience any of the following mild to moderate symptoms.** High blood sugar levels over a long period of time can lead to too much acid in the blood (diabetic ketoacidosis).

Symptoms of mild to moderate hyperglycaemia may include:

- drowsiness
- flushed face
- thirst, loss of appetite
- fruity odour on the breath
- blurred vision
- passing larger amounts of urine than usual
- getting up at night more often than usual to pass urine
- high levels of glucose and acetone in the urine
- nausea and vomiting.

**Contact your doctor or the Emergency Department immediately if your blood sugar level is too high and you experience any of the following severe symptoms.** Severe hyperglycaemia can lead to unconsciousness and in extreme cases death (if left untreated).

Symptoms of severe hyperglycaemia may include:

- heavy breathing
- fast pulse
- nausea, vomiting
- nausea and vomiting where you cannot keep any fluids down (leading to severe dehydration)
- loss of consciousness.

### **Illness**

**Tell your doctor if you are ill.**

Illness, especially with nausea and vomiting, may cause your insulin needs to change. Even if you are not eating, you still require insulin. You and your doctor should design an insulin plan for those times when you are sick.

### **Emotional disturbances**

**Tell your doctor if you are feeling particularly excited, upset, or stressed.**

Emotional disturbances may impact your dose of HUMULIN.

### **Exercise**

Exercise may lower your need for insulin. Exercise may also make insulin work more quickly, especially if the exercise involves the area around the injection site (e.g., the leg should not be used as an injection site prior to jogging or running).

**Tell your doctor if you are exercising more than usual, as your dose of HUMULIN may need to change.**

## Diet

Changes to your diet may cause your insulin needs to change.

**Tell your doctor before making any significant changes to your diet (e.g., removing carbohydrates).**

## Injection site

If you inject HUMULIN in the same place, the fatty tissue may shrink (lipoatrophy) or thicken (lipohypertrophy).

Lumps under the skin may also be caused by build-up of a protein called amyloid (cutaneous amyloidosis).

To help prevent these skin changes, **do not use the same place for injection more often than once a month.**

HUMULIN may not work very well if you inject into a lumpy, shrunken, or thickened area. **Avoid injecting into these areas.**

**Tell your doctor if you are currently injecting into these affected areas before you start injecting in a different area.**

Your doctor may tell you to measure your blood sugar more frequently, and to adjust your dose of HUMULIN or other antidiabetic medications.

## Driving or using machines

**Be careful before you drive or use any machine or tools until you know how HUMULIN affects you.**

**Do not drive if you feel like your blood sugar level is low.**

Hypoglycaemia (low blood sugar) may impair your ability to concentrate or react while driving.

## Drinking alcohol

**Tell your doctor if you drink alcohol.**

Alcohol may mask the symptoms of hypoglycaemia (low blood sugar) and make it more difficult to recognise them.

Alcohol may also affect the dose of HUMULIN that you are prescribed.

## Looking after your medicine

- Keep your cartridges in a fridge between 2°C - 8°C. Do not freeze.
- You may keep the cartridges below 30°C for up to 21 days while you are using them.
- When the cartridge has been inserted into the pen, the cartridge pen combination should not be kept in the fridge.
- Dispose of unrefrigerated cartridges after 21 days even if they still contain insulin.
- Do not store the cartridges in the bathroom or near a sink.
- Do not put the cartridges near heat or in the sun.
- **Keep it where young children cannot reach it.**

## When to discard your medicine

Discard cartridges after the expiry date (month, year) stamped on the label has passed.

## Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Empty cartridges and needles should be disposed of in a sharps container or similar puncture proof container composed of hard plastic or glass.

**Ask your doctor, nurse, pharmacist or diabetes educator where you can dispose of the container once it is full.**

## 6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

### Less serious side effects

Less serious side effects	What to do
<b>Blood sugar related:</b> <ul style="list-style-type: none"><li>• <b>Mild to moderate hypoglycaemia</b> (low blood sugar), symptoms may include:<ul style="list-style-type: none"><li>o weakness, trembling or shaking</li><li>o sweating</li><li>o light headedness/headache</li><li>o lack of concentration</li><li>o behaviour change</li><li>o dizziness</li><li>o tearfulness/crying</li><li>o irritability</li><li>o numbness around the lips/fingers</li><li>o hunger.</li></ul></li></ul>	<b>Immediately eat some sugary food or drink, such as jelly beans, fruit juice, or glucose tablets</b> <b>If you do not feel better after eating/drinking some sugary food or drink, contact your doctor or go to the Emergency Department at your nearest hospital.</b>
<b>Allergy related:</b> <ul style="list-style-type: none"><li>• <b>Local allergy</b>, symptoms may include redness, swelling and itching around the injection site. These symptoms generally disappear within a few weeks of continued use.</li></ul> <b>Skin related:</b> <ul style="list-style-type: none"><li>• <b>Lipodystrophy</b>, symptoms may include a depression in the skin or an enlargement or thickening of the tissue around the injection site.<ul style="list-style-type: none"><li>o A change in injection technique may help resolve the problem.</li></ul></li></ul>	<b>Speak to your doctor if you have any of these less serious side effects and they worry you.</b>

Less serious side effects	What to do
<b>General:</b> <ul style="list-style-type: none"> <li>• <b>Oedema (fluid retention)</b>, particularly in the ankles, feet and hands.</li> </ul>	

### Serious side effects

Serious side effects	What to do
<b>Blood sugar related:</b> <ul style="list-style-type: none"> <li>• <b>Severe hypoglycaemia</b> (low blood sugar), symptoms may include: <ul style="list-style-type: none"> <li>o disorientation</li> <li>o seizures, fits, or convulsions</li> <li>o loss of consciousness.</li> </ul> </li> </ul>	<p><b>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</b></p> <p><b>If you lose consciousness, make sure friends and family know to turn you on your side and get medical help immediately. They should <u>not</u> give you anything to eat or drink to avoid choking.</b></p>
<b>Allergy related:</b> <ul style="list-style-type: none"> <li>• <b>Generalised allergy</b>, symptoms may include <ul style="list-style-type: none"> <li>o rash over a large part of the body</li> <li>o shortness of breath</li> <li>o wheezing</li> <li>o low blood pressure</li> <li>o swelling of the face, lips, or tongue</li> <li>o fast pulse</li> <li>o sweating</li> </ul> </li> </ul>	<p><b>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</b></p>

**Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.**

Other side effects not listed here may occur in some people.

### Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems). By reporting side

effects, you can help provide more information on the safety of this medicine.

**Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.**

## 7. Product details

HUMULIN is only available with a doctor's prescription.

### What HUMULIN contains

<b>Active ingredient (main ingredient)</b>	biosynthetic human insulin (100 units per mL)
<b>Other ingredients (inactive ingredients)</b>	<p><b>HUMULIN R</b></p> <ul style="list-style-type: none"> <li>• metacresol</li> <li>• glycerol</li> <li>• hydrochloric acid</li> <li>• sodium hydroxide</li> <li>• water for injections</li> </ul> <p><b>HUMULIN NPH and HUMULIN 30/70</b></p> <ul style="list-style-type: none"> <li>• metacresol</li> <li>• glycerol</li> <li>• zinc oxide</li> <li>• dibasic sodium phosphate heptahydrate</li> <li>• protamine sulfate</li> <li>• phenol</li> <li>• hydrochloric acid</li> <li>• sodium hydroxide</li> <li>• water for injections</li> </ul>

**Do not take this medicine if you are allergic to any of these ingredients.**

### What HUMULIN looks like

#### HUMULIN R (AUST R 39666)

HUMULIN R is available in 3 mL cartridges.

HUMULIN R is a clear colourless solution with a water-like appearance and consistency.

#### HUMULIN NPH (AUST R 39662)

HUMULIN NPH is available in 3 mL cartridges.

HUMULIN NPH is a white and cloudy or milky suspension.

#### HUMULIN 30/70 (AUST R 39659)

HUMULIN 30/70 is available in 3 mL cartridges.

HUMULIN 30/70 cartridges is a white and cloudy or milky suspension.

### Who distributes HUMULIN?

#### Supplied in Australia by:

Eli Lilly Australia Pty Ltd

Level 9, 60 Margaret Street

SYDNEY NSW 2000

**Supplied in New Zealand by:**

Eli Lilly and Company (NZ) Limited

PO Box 109 197

Newmarket

Auckland 1149

® = Registered Trademark

This leaflet was prepared in October 2024.

vA6.0\_Oct2024