

Minims[®] Oxybuprocaine

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using Minims[®] Oxybuprocaine?

Minims[®] Oxybuprocaine contains the active ingredient oxybuprocaine hydrochloride. Minims[®] Oxybuprocaine is used to numb or block the feeling of pain in your eye(s) during short-term medical procedures. These procedures may include: measuring the pressure in your eye, inserting contact lenses, removing foreign bodies from your eye(s) and other minor operations on your eye(s).

For more information, see Section [1. Why am I using Minims[®] Oxybuprocaine?](#) in the full CMI.

2. What should I know before I use Minims[®] Oxybuprocaine?

Do not use if you are allergic to oxybuprocaine, any other local anaesthetic or any of the ingredients listed at the end of this CMI. Do not use if you have an infection

in your eye. **Talk to your doctor if you have any other medical conditions or take any other medicines.**

For more information, see Section [2. What should I know before I use Minims® Oxybuprocaine?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with Minims® Oxybuprocaine and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use Minims® Oxybuprocaine?

- Your doctor will instill Minims® Oxybuprocaine eye drops into your eye(s) before starting a procedure on your eye(s).
- The usual dose is one drop instilled into the eye.
- The number of times your doctor will put drops in your eye(s) will depend on the type and length of the procedure to be performed.

More instructions can be found in Section [4. How do I use Minims® Oxybuprocaine?](#) in the full CMI.

5. What should I know while using Minims® Oxybuprocaine?

Things you should do	<ul style="list-style-type: none">• Protect your numb eye from dust particles,
-----------------------------	--

	which could cause infections.
Things you should not do	<ul style="list-style-type: none"> • Do not touch or rub your eye while your eye is numb.
Driving or using machines	<ul style="list-style-type: none"> • Be careful before you drive or use any machines or tools until you know how Minims® Oxybuprocaine affects you. • Keep the patch that your doctor puts on your eye until normal feeling returns. • Do not drive or operate hazardous machinery until you regain clear vision.
Looking after your medicine	<ul style="list-style-type: none"> • Your doctor will keep Minims® Oxybuprocaine in a refrigerator (2-8°C). The eye drops should not be frozen or exposed to strong light.

For more information, see Section [5. What should I know while using Minims® Oxybuprocaine?](#) in the full CMI.

6. Are there any side effects?

All medicines can have side effects. If they do occur, they are usually minor and temporary. Do not be alarmed by this list. You may not experience any of them.

Common side effects include: stinging or burning sensation in the eye. Serious side effects that require urgent medical attention include: ring-like structure in the outer part of the cornea, cardiovascular and central nervous system symptoms and symptoms of an allergic reaction.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

Minims® Oxybuprocaine

Active ingredient: *Oxybuprocaine hydrochloride*

Consumer Medicine Information (CMI)

This leaflet provides important information about using Minims® Oxybuprocaine. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Minims® Oxybuprocaine.**

Where to find information in this leaflet:

- [1. Why am I using Minims® Oxybuprocaine?](#)
- [2. What should I know before I use Minims® Oxybuprocaine?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use Minims® Oxybuprocaine?](#)
- [5. What should I know while using Minims® Oxybuprocaine?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using Minims® Oxybuprocaine?

Minims® Oxybuprocaine contains the active ingredient oxybuprocaine hydrochloride. Minims® Oxybuprocaine belongs to a class of medicines called local anaesthetics.

Minims® Oxybuprocaine is used to numb or block the feeling of pain in your eye(s) during short-term medical procedures. These procedures may include: measuring the pressure in your eye, inserting contact lenses, removing foreign bodies from your eye(s) and other minor operations on your eye(s).

2. What should I know before I use Minims® Oxybuprocaine?

Warnings

Do not use Minims® Oxybuprocaine if:

- You are allergic to oxybuprocaine, any other local anaesthetic or any of the ingredients listed at the end of this leaflet. Always check the ingredients to make sure you can use this medicine.
- You have an infection in your eye.

Check with your doctor if:

- You have had an allergy to any other medicines, foods, preservatives or dyes.
- You have any other medical conditions.
- You take any medicines for any other condition.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

This medicine has been developed for use in the eye only. It should not be swallowed or injected.

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Minims® Oxybuprocaine may interfere with each other. These medicines may be affected by Minims® Oxybuprocaine or may affect how well it works. These include:

- A group of medicines called anticholinesterases
- A medicine called suxamethonium

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Minims® Oxybuprocaine.

4. How do I use Minims® Oxybuprocaine?

Your doctor will not give you a prescription for your own use. This medicine has been specifically developed for use by your doctor to numb the feeling in your eye(s) while he/she is attending to your eye(s), and not for long-term use.

How much to use

- The usual dose is one drop into your eye(s) before starting a procedure on your eye(s).
- For most procedures one or two drops will be sufficient. For the removal of foreign bodies or minor surgery, three to six drops may be required.
- The number of times your doctor will put drops in your eye(s) will depend on the type and length of the procedure to be performed.

How to use Minims® Oxybuprocaine

- Your doctor will put the correct number of drops of Minims® Oxybuprocaine into your eye(s) before starting a procedure on your eye(s).
- If you are wearing contact lenses, you will be asked to remove them before putting in the eye drops. You may put your contact lenses back into your eye after the numbness has worn off or when your doctor tells you to.
- In children, the doctor may compress the small part of the child's eye near the corner of his/her nose for a minute during and following the instillation of the drops.

If you use too much Minims® Oxybuprocaine

Minims® Oxybuprocaine can damage the surface of the eye or cause tolerance (when your body becomes less responsive to the effects of a medicine over time, requiring higher doses to achieve the same effect) if it is used too often.

The doctor giving you Minims® Oxybuprocaine will be experienced in its use, so it is unlikely that you will be given an overdose.

If too many drops have accidentally been put into your eye(s):

- Immediately rinse your eye(s) with warm water (if you are wearing contact lenses, remove them before the rinse).

Accidental swallowing of the solution is unlikely to cause any bad side effects due to the low content of oxybuprocaine.

If you think that you or anyone else may have swallowed any or all of the contents of the container of Minims® Oxybuprocaine, you should immediately:

- phone the Poisons Information Centre (in Australia, call **13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while using Minims® Oxybuprocaine?

Things you should do

- Protect your numb eye from dust particles, which could cause infections.

Things you should not do

- Do not touch or rub your eye while your eye is numb.

Your eye(s) may remain numb for up to one hour after the last drop is put in.

Your doctor will put a patch over your eye after finishing the procedure to protect your eye from dust and bacterial contamination and stop you from touching it, scratching the surface of the eye and damaging it.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how Minims® Oxybuprocaine affects you.

Keep the patch that your doctor puts on your eye until normal feeling returns. Do not drive or operate hazardous machinery until you regain clear vision.

Looking after your medicine

Your doctor will keep Minims® Oxybuprocaine in a refrigerator (2-8°C). The eye drops should not be frozen or exposed to strong light.

When to discard your medicine

Your doctor will discard each single dose unit and any remaining solution in it immediately after use. He/she will not save unused contents.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<p>Eyes:</p> <ul style="list-style-type: none">• Stinging or burning sensation• Blurred vision and/or problems seeing clearly• Local irritation or feeling of having something in the eye• Redness• Itching• Swelling• Dry eye <p>Skin:</p> <ul style="list-style-type: none">• Rash <p>Gastrointestinal symptoms, such as:</p> <ul style="list-style-type: none">• Nausea and vomiting• Difficulty swallowing food, liquids or saliva	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Serious side effects

Serious side effects	What to do
<p>Eyes:</p> <ul style="list-style-type: none">• Ring-like structure in the outer part of the cornea <p>Cardiovascular symptoms, such as:</p> <ul style="list-style-type: none">• Slow heartbeat <p>Central nervous system symptoms, such as:</p> <ul style="list-style-type: none">• Drowsiness• Confusion or disorientation• Agitation or feeling of extreme happiness• Visual, hearing or speech disorders• Tingling sensation in the skin• Muscle twitching• Seizure <p>Symptoms of an allergic reaction, such as:</p> <ul style="list-style-type: none">• Shortness of breath	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

Serious side effects	What to do
<ul style="list-style-type: none"> ● Wheezing or difficulty breathing ● Swelling of the face, lips, tongue or other parts of the body ● Rash, itching or hives on the skin 	

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects online (in Australia, go to www.tga.gov.au/reporting-problems). By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

What Minims[®] Oxybuprocaine contains

Active ingredient (main ingredient)	oxybuprocaine hydrochloride (0.4% w/v)
Other ingredients (inactive ingredients)	<ul style="list-style-type: none">• Purified water• Hydrochloric acid

Do not take this medicine if you are allergic to any of these ingredients.

The eye drops do not contain any preservatives.

What Minims[®] Oxybuprocaine looks like

Minims[®] Oxybuprocaine comes in single-dose units (tubes) containing 0.5 mL of a clear, colourless solution. Each tube is wrapped in a polyester/paper blister. The blisters are packed in cartons of 20 units (AUST R 32259).

Who distributes Minims[®] Oxybuprocaine

In Australia, Minims[®] Oxybuprocaine is supplied by:

Bausch & Lomb (Australia) Pty Ltd

Level 2, 12 Help Street

Chatswood, NSW 2067

Phone: 1800 251 150

This leaflet was prepared in January 2025.