

Prolia[®]

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using Prolia?

Prolia contains the active ingredient denosumab. Prolia is used to improve bone density and to reduce your risk of fracture. It is used to treat bone loss in women with osteoporosis after the menopause, men with osteoporosis, and men with prostate cancer who have reduced testosterone level caused by surgery or treatment with drugs. It is also used to improve bone density in patients treated with corticosteroids.

For more information, see Section [1. Why am I using Prolia?](#) in the full CMI.

2. What should I know before I receive Prolia?

Do not use if you have ever had an allergic reaction to denosumab, medicines produced using Chinese Hamster Ovary cells or any of the ingredients listed at the end of the CMI.

Do not use in patients under 18 years of age.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

Tell your doctor if you have calcium deficiency.

For more information, see Section [2. What should I know before I use Prolia?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with Prolia and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use Prolia?

The recommended dose of Prolia is 60 mg given once every 6 months as a single injection under the skin.

More instructions can be found in Section [4. How do I use Prolia?](#) in the full CMI.

5. What should I know while using Prolia?

Things you should do	
	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist

	<p>you visit that you are using Prolia.</p> <ul style="list-style-type: none"> ● Take calcium and vitamin D supplements if your doctor has told you to. ● Maintain good oral hygiene when being treated with Prolia. ● Attend all of your treatment and doctor's appointments so that your progress can be checked. ● Tell your doctor immediately if you become pregnant while taking Prolia.
<p>Things you should not do</p>	<ul style="list-style-type: none"> ● Do not stop using Prolia without talking to your doctor.
<p>Looking after your medicine</p>	<ul style="list-style-type: none"> ● Store Prolia in the refrigerator (2°C to 8°C) in the original pack to protect from light. Do not freeze. ● Do not shake or vigorously agitate the pre-filled syringe.

For more information, see Section [5. What should I know while using Prolia?](#) in the full CMI.

6. Are there any side effects?

Side effects that require urgent medical attention include: signs of an allergic reaction; muscle aches, twitches or cramps; numbness or tingling in your fingers, toes or around your mouth; persistent pain or swelling and/or non-healing sores in your mouth or jaw; pain in your hip, groin, or thigh, which is sometimes severe; severe allergic reaction with skin rash, blisters or fever.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

Prolia[®] (den"-os"-u"-mab)

Active ingredient(s): *denosumab*

Consumer Medicine Information (CMI)

This leaflet provides important information about using Prolia. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Prolia.**

Where to find information in this leaflet:

- [1. Why am I using Prolia?](#)
- [2. What should I know before I use Prolia?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use Prolia?](#)
- [5. What should I know while using Prolia?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using Prolia?

Prolia contains the active ingredient denosumab. Denosumab is a protein (monoclonal antibody) that attaches (binds) specifically to another unique protein in the body in order to stop the development of bone-removing cells before they reach the bones and cause

damage. Treatment with Prolia makes your bone stronger and less likely to break.

Prolia is used to:

- Treat osteoporosis in women after the menopause, to reduce the risk of spinal, non-spinal and hip fractures.
- Treat bone loss in men with osteoporosis at increased risk of fracture.
- Treat bone loss that results from a reduction in testosterone level caused by surgery or treatment with drugs in men with prostate cancer.
- Improve bone density in patients treated with corticosteroids.

Bone is a living tissue and is renewed all the time. In women, the ovaries produce the hormone oestrogen which helps keep bones healthy. After menopause, the oestrogen level drops which affects the bone renewal cycle so that more bone is lost than made, resulting in a lower bone mass. This leaves bones thin and fragile. Osteoporosis is the term used to describe an increased fracture risk, usually with low bone density.

Osteoporosis becomes more common with increasing age. It is more common in women. It can also occur in patients receiving corticosteroids. Many people with osteoporosis have no symptoms but they are still at risk of breaking bones (developing fractures), especially in the spine, hips and wrists.

Other things that can increase the risk of fractures include:

- age

- existence of a previous fracture
- family history of hip fractures
- low body weight
- drinking alcohol
- smoking.

Prolia is prescribed to improve your bone density and to reduce your risk of fracture.

Surgery or medicines used in the treatment of men with prostate cancer to stop the production of testosterone can also lead to bone loss. The bones become weaker and break more easily.

Your doctor, however, may have prescribed Prolia for another reason.

2. What should I know before I use Prolia?

Warnings

Do not use Prolia if:

- you have low calcium levels in your blood (hypocalcaemia). Your doctor may do a blood test to check your calcium levels before you use Prolia.
- you are pregnant, think you may be pregnant, or trying to get pregnant. Prolia may harm your unborn baby.
- you are breast-feeding. It is not known if the active ingredient, denosumab, passes into breast milk.

- you are allergic to denosumab, any medicines that are produced using Chinese Hamster Ovary cells, or any of the ingredients listed at the end of this leaflet. Always check the ingredients to make sure you can use this medicine.
- you are a child or adolescent. Prolia is not indicated for use in patients under 18 years of age.
- you are taking another medicine containing denosumab.
- the expiry date [EXP.] printed on the pack has passed. If you use it after the expiry date has passed, it may not work as well.
- the packaging is torn or shows signs of tampering.
- the Prolia solution is cloudy or discoloured. There may be some translucent to white particles of protein in the solution, however the medicine can still be used.

Check with your doctor if you:

- have allergies to any other medicines, or any other substances such as foods, preservatives or dyes.
- have calcium or vitamin D deficiency. If you are prone to low calcium levels, your doctor will monitor your blood especially in the first few weeks after starting Prolia. Severe low blood calcium levels may lead to hospitalisation, life-threatening events and death.
- are unable to take daily calcium or vitamin D supplements.
- have or have had severe kidney problems, kidney failure or have needed dialysis, which may increase

your chance of getting low blood calcium if you do not take calcium supplements.

- had or have pain in the teeth, gums or jaw, swelling or numbness of the jaw, a "heavy jaw feeling" or loosening of a tooth. A dental condition called jaw osteonecrosis has been rarely reported in patients treated with Prolia. Your doctor should examine your mouth and may ask for a dental examination before you start Prolia. You may need to have dental treatment completed before starting your medicine. You may have higher risk of developing jaw problems if you:
 - are undergoing chemotherapy, taking steroids, or are having a dental procedure
 - do not receive routine dental care, have gum disease or have taken Prolia for a long time
- have been told by a doctor or other healthcare professional that you have an intolerance to some sugars.
- take any medicines for any other condition.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy

Prolia has not been tested in pregnant women. Do not use Prolia if you are pregnant, think you may be pregnant, or trying to get pregnant. Prolia may harm your unborn baby.

Breastfeeding

Do not use Prolia if you are breast-feeding. It is not known if the active ingredient, denosumab, passes into breast milk. It is important to talk to your doctor if you are breast-feeding or plan to do so. Your doctor will then help you decide whether to stop breast-feeding or whether to stop taking Prolia.

Use in children

Do not use Prolia in children or adolescents under 18 years of age.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Tell your doctor if you are taking another medicine containing denosumab (Xgeva). If you are taking Xgeva, and/or other medicines containing denosumab, you should not take Prolia.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Prolia.

4. How will I be given Prolia?

How is Prolia given

- Prolia is given as an injection under the skin. This is called a subcutaneous injection.

How much Prolia will I be given

- For each dose of Prolia you will be given 60 mg as a single injection.

When you will be given Prolia

- Prolia is injected once every 6 months.
- It is important to take your dose of Prolia on schedule. Do not skip or delay getting your injection.
- Each pack of Prolia contains a reminder card that can be removed from the carton. Use the reminder card to keep a record of the next injection date.
- Continue using Prolia for as long as your doctor tells you to. Prolia can treat osteoporosis and bone loss only for as long as you keep having treatment. Please talk to your doctor before you consider stopping treatment.
- You should also take calcium and vitamin D supplements while receiving Prolia. Prolia may lower the calcium levels in your blood. Your doctor, nurse or pharmacist will discuss how much calcium and vitamin D you should take to help prevent low calcium levels.


Instructions for injecting Prolia when supplied in a pre-filled syringe with an automatic needle guard

This section contains information on how you, or the person injecting you (your carer), must use the Prolia pre-filled syringe.

To reduce the risk of accidental injury by the needle, each pre-filled syringe is equipped with an automatic needle guard that is automatically activated to cover the needle after complete delivery of the pre-filled syringe contents.

Guide to parts

Before and after use



Important

Before you or your carer use Prolia pre-filled syringe with automatic needle guard, read this important information:

- It is important that you or your carer do not try to give the injection unless training from a doctor, nurse or pharmacist has been received.

- Prolia is given as an injection into the tissue just under the skin (subcutaneous injection).
- DO NOT remove the grey needle cap from the pre-filled syringe until you are ready to inject.
- DO NOT use the pre-filled syringe if it has been dropped on a hard surface. Use a new pre-filled syringe and call your doctor, nurse or pharmacist.
- DO NOT attempt to activate the pre-filled syringe prior to injection.
- DO NOT attempt to remove the clear pre-filled syringe safety guard from the pre-filled syringe.

Call your doctor, nurse or pharmacist if you have any questions.

Step 1: Prepare

A: Remove the pre-filled syringe tray from the package and gather the supplies needed for your injection: alcohol wipes, a cotton ball or gauze pad, a plaster and a sharps disposal container (not included).

For a more comfortable injection, leave the pre-filled syringe at room temperature for about 30 minutes before injecting. Wash your hands thoroughly with soap and water.


On a clean, well-lit work surface, place the new pre-filled syringe and the other supplies.

- DO NOT try to warm the syringe by using a heat source such as hot water or microwave.
- DO NOT leave the pre-filled syringe exposed to direct sunlight.

- DO NOT shake the pre-filled syringe.

Keep pre-filled syringes out of the sight and reach of children.


B: Open the tray, peeling away the cover. Grab the pre-filled syringe safety guard to remove the pre-filled syringe from the tray.



For safety reasons:

- DO NOT grasp the plunger.
- DO NOT grasp the grey needle cap.

C: Inspect the medicine and pre-filled syringe




DO NOT use the pre-filled syringe if:

- The medicine is cloudy or there are particles in it. It must be a clear, colourless to slightly yellow solution.
- Any part appears cracked or broken.
- The grey needle cap is missing or not securely attached.
- The expiry date printed on the label has passed the last day of the month shown.

In all cases, call your doctor, nurse or pharmacist.

Step 2: Get ready

A: Wash your hands thoroughly. Prepare and clean your injection site.




You can use:

- Upper part of your thigh.
- Belly, except for a 5 cm (2-inch) area right around your belly button.
- Outer area of upper arm (only if someone else is giving you the injection).


Clean the injection site with an alcohol wipe. Let the skin dry.

- DO NOT touch the injection site before injecting.
- DO NOT inject into areas where the skin is tender, bruised, red, or hard. Avoid injecting into areas with scars or stretch marks.

B: Carefully pull the grey needle cap straight out and away from your body.



C: Pinch the injection site to create a firm surface.



It is important to keep the skin pinched when injecting.


Step 3: Inject

A: Hold the pinch. INSERT the needle into skin.




- DO NOT touch the cleaned area of the skin.

B: PUSH the plunger with slow and constant pressure until you feel or hear a "snap". Push all the way down through the snap.



It is important to push down through the "snap" to deliver your full dose.

C: RELEASE your thumb. Then LIFT the syringe off the skin.




After releasing the plunger, the pre-filled syringe safety guard will safely cover the injection needle.

- DO NOT put the grey needle cap back on used pre-filled syringes.

Step 4: Finish

A: Discard the used pre-filled syringe and other supplies in a sharps disposal container.



Medicines should be disposed of in accordance with local requirements. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Keep the syringe and sharps disposal container out of sight and reach of children.

- DO NOT reuse the pre-filled syringe.
- DO NOT recycle pre-filled syringes or throw them into household waste.

B: Examine the injection site

If there is blood, press a cotton ball or gauze pad on your injection site. DO NOT rub the injection site. Apply a plaster if needed.

If you miss a dose of Prolia

If you miss a dose, Prolia should be administered as soon as possible. From then on, Prolia should be scheduled every 6 months from the date of the last injection.

If you are given too much Prolia

If you think you or anyone else has received too much Prolia, you should immediately:

- phone the Poisons Information Centre (**by calling 13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while using Prolia?

Things you should do

- take calcium and vitamin D supplements if your doctor has told you to. Most people do not get enough calcium and vitamin D in their diet and supplements are needed to help strengthen bones.
- practice good dental hygiene while being treated with Prolia.

Your routine dental hygiene should include brushing your teeth and tongue after every meal, including the evening and gentle flossing once a day to remove plaque.

Use a mirror and check your teeth and gums regularly for any changes such as sores or bleeding gums. If you notice any problems, tell your doctor and dentist immediately.

- tell any doctor, dentist, nurse, or pharmacist you visit that you are using Prolia.
- if you are about to be started on any other medicine, remind your doctor, nurse or pharmacist that you are being treated with Prolia.
- attend all your doctor's appointments so that your progress can be checked. Your doctor may recommend you to have some tests, X-rays and/or bone density scans from time to time to make sure the medicine is working.

Call your doctor straight away if you:

- have spasms, twitches, aches or cramps in your muscles, and/or numbness or tingling in your fingers, toes or around your mouth, or have seizures. You may have low levels of calcium in your blood.
- experience any problems with your mouth or teeth such as loose teeth or ill-fitting dentures, pain, or swelling while being treated with Prolia.
- develop a swollen, red area of skin, most commonly in the lower leg, that feels hot and tender (cellulitis) and sometimes experienced with fever and chills.

- experience new or unusual pain in your hip, groin, or thigh. Some people have developed unusual fractures in their thigh bone while being treated with Prolia. These fractures may occur with little or no trauma and may involve both thigh bones. This side effect is very rare.
- experience shortness of breath; wheezing or difficulty breathing; swelling of the face, lips, tongue, throat or other parts of the body; rash, itching or hives on the skin. If you experience any of these side effects you may be having an allergic reaction to Prolia. These side effects are rare.
- have a severe allergic reaction with skin rash, blisters or fever
- notice any purple or brownish-red spots, hives or skin sores. This may be an allergic reaction that can damage blood vessels mainly in the skin. This side effect is very rare.
- become pregnant while using Prolia. Your doctor can discuss with you the risks of having it while you are pregnant.
Females that are menstruating should ensure they have adequate contraception while taking Prolia.

Things you should not do

- Do not stop using Prolia without talking to your doctor. After your treatment with Prolia is stopped, or if you skip or delay taking a dose, your risk of breaking bones in your spine is increased, especially if you have a history of broken bones in the spine. If your

Prolia treatment is stopped, discuss other available treatment options with your doctor.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how Prolia affects you.

Prolia has no known effects on the ability to drive or use machines but, as a general precaution, if you are driving soon after an injection, arrange to have someone else drive.

Looking after your medicine

If you need to store your Prolia before use, follow the instructions in the carton on how to take care of your medicine properly.

- Store Prolia in the refrigerator between 2 and 8°C. Do not freeze.
- Keep your medicine in the original carton to protect from light. If you remove the medicine from the carton it may not keep well.
- Your medicine may be left outside the refrigerator to reach room temperature (up to 25°C) before injection. This will make the injection more comfortable.
- Once your medicine has been left to reach room temperature (up to 25°C), it must be used within 30 days.
- Do not keep Prolia at temperatures above 25°C. Warm temperatures will affect how Prolia works.

- Do not shake or vigorously agitate the pre-filled syringe.

Keep it where young children cannot reach it.

When to discard your medicine

Prolia is for single-use in one patient only. Dispose of any unused or expired medicine as instructed below.

Getting rid of any unwanted medicine

Your doctor or nurse is likely to dispose of Prolia for you. However, if you need to get rid of this medicine because you no longer need to use it or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<p>Muscle and skeleton:</p> <ul style="list-style-type: none">• back, muscle or joint pain, or stiffness, most commonly affecting the hips, knees and spine• pain in the arms or legs• aching muscle, muscle tenderness or weakness, not caused by exercise <p>Skin and hair:</p> <ul style="list-style-type: none">• unusual hair loss or thinning• itchy, red or dry skin <p>Ears:</p> <ul style="list-style-type: none">• ear pain, discharge from the ear and/or an ear infection. These could be signs of bone damage in the ear <p>Eyes:</p> <ul style="list-style-type: none">• blurred or cloudy vision <p>General:</p>	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Less serious side effects	What to do
<ul style="list-style-type: none"> • high cholesterol levels in the blood 	

Serious side effects

Serious side effects	What to do
<p>Signs of an allergic reaction:</p> <ul style="list-style-type: none"> • shortness of breath • wheezing or difficulty breathing. • swelling of the face, lips, tongue or other parts of the body • rash, itching or hives on the skin. • rash that may occur on the skin or sores in the mouth <p>Signs of low blood calcium (hypocalcaemia):</p> <ul style="list-style-type: none"> • muscle spasms, twitches, aches or cramps 	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

Serious side effects	What to do
<ul style="list-style-type: none"> ● numbness or tingling in your fingers, toes or around your mouth ● seizures <p>Signs of problems with your mouth, teeth or jaw:</p> <ul style="list-style-type: none"> ● persistent pain or swelling and/or non-healing sores in your mouth or jaw ● loose teeth <p>Signs of bone fractures:</p> <ul style="list-style-type: none"> ● bone, joint and/or muscle pain, including pain in your hip, groin, or thigh, which is sometimes severe ● pain in spine <p>Signs of skin infection:</p> <ul style="list-style-type: none"> ● develop a swollen, red area of skin that feels hot and tender (cellulitis) and sometimes experienced with fever and chills ● skin rash or blisters with fever 	

Serious side effects	What to do
<p>Gut and digestion:</p> <ul style="list-style-type: none"> • pain in upper abdomen (belly) that may be accompanied by back pain, nausea, vomiting, fever and/or sweating <p>Brain and nerves:</p> <ul style="list-style-type: none"> • pain in the extremities, such as hands and feet 	

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What Prolia contains

Active ingredient (main ingredient)	denosumab
Other ingredients (inactive ingredients)	Acetate sodium hydroxide sorbitol polysorbate 20 water for injections

Do not take this medicine if you are allergic to any of these ingredients.

What Prolia looks like

Prolia is a clear, colourless to slightly yellow solution supplied in a pre-filled syringe with an automatic needle guard.

Prolia comes in a single pack size containing 60 mg of denosumab in a volume of 1.0 mL (60 mg/1.0 mL). Aust R 159323.

Each pack contains one ready to use single-use pre-filled syringe and one reminder card.

The needle cover on the pre-filled syringe is not made with natural rubber latex.

Who distributes Prolia

Amgen Australia Pty Ltd

Level 11, 10 Carrington St

Sydney NSW 2000

Ph: 1800 803 638

www.amgenmedinfo.com.au

This leaflet was prepared in March 2025.